SECI	IRITY	CLASSIFICATION OF	THIS	PAGE

SECURITY CLASSIFICATION OF THIS PAGE						
	REPORT DOCU	MENTATION	PAGE			
1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE	MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY	3 DISTRIBUTION/AVAILABILITY OF REPORT					
2b. DECLASSIFICATION / DOWNGRADING SCHED	Approved for public release; distribution is unlimited					
4 PERFORMING ORGANIZATION REPORT NUMB	5. MONITORING ORGANIZATION REPORT NUMBER(S)					
CR 88-09		CR 88-	09			
6a. NAME OF PERFORMING ORGANIZATION Science Applications International Corp.	6b OFFICE SYMBOL (If applicable)		ONITORING ORG Environmen Research F	tal Predi	ction	
6c. ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (Ci	ty, State, and ZII	Code)		
205 Montecito Ave. Monterey, CA 93940		Monte	rey, CA 939	43-5006		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Commander, Naval Oceanography Command	8b. OFFICE SYMBOL (If applicable)		it instrument ii 8-84-D-3187		NUMBER	
8c. ADDRESS (City, State, and ZIP Code)	<u> </u>	10 SOURCE OF	FUNDING NUMBE	RS		
NSTL, MS 39529-5000		PROGRAM PROJECT TASK NO.			WORK UNIT ACCESSION NO.	
Funding s To COSATI CODES FIELD GROUP SUB-GROUP 19 ABSTRACT (Continue on reverse if necessary This handbook for the pofor Mediterranean ports, proviare threatened by actual or for thunderstorms in the port vicidiscussed. Precautionary or entered the polymer of the handbook content and use; a question of the polymer.	Source: 0&M,N-1 18. SUBJECT TERMS (C. Storm haven La Spezia por and identify by block of the control of La Spezia, ides decision-malorecast strong winity. Causes are evasive actions for ick-look captain	Continue on revers t comber) one in a seking guidance inds, high sind effects of the suggester ready refencies suggester ready refencies suggester.	e if necessary and Mediterr Mediterr Mediterr Mediterr for ship eas, restrif such hazad for variorence: gene	d identify by anean meta anean ocea vere weath captains worked visibly rdous concus vessel ral guidanailed rev	block number) eorology anography er guides whose vessels oility or ditions are situations. nce on iew of	
general information on environ oceanographic information. 20 DISTRIBUTION/AVAILABILITY OF ABSTRACT SUNCLASSIFIED/UNLIMITED SAME AS 223 NAME OF RESPONSIBLE INDIVIDUAL Perryman, Dennis C., contract	RPT DTIC USERS	21. ABSTRACT SE	CURITY CLASSIFIC	CATION D	E SYMBOL	
DD FORM 1472 84 MAR 87 A		(400) 047	1703	ן ייטאין	917 - 1	

```
AD-4198 909
        040200
FG (2)
        080300
CI (3)
        (11)
        SCIENCE APPLICATIONS INTERNATIONAL CORP MONIEREY CA
CA (5)
        Severe Weather Guide . Mediterranean Ports. 21. La
T1(6)
        Sperla.
TC (8)
        (11)
        Final rept. 13 Sep 84-1 Nov 86.
DN (9)
All (10) Englebretson, Ronald E.
AU (10)
        Gilmore, Richard D.
RD (11) May 1988
PG (12) 64
CT (15) NØØ228-84-D-3137
RN (18) NEPRF-CR-88-09
RC (20) Unclassified report
        *LIGURIAN SEA. *MARINE METEOROLOGY, *HARBORS, *SAFETY, *
DE (23)
        STORMS, DECISION MAKING, HANDBOOKS, HAZARDS, ITALY.
        LIMITATIONS. MEDITERRANEAN SEA. METEOROLOGICAL DATA.
        OCEAN WAVES, OCEANOGRAPHIC DATA, PORTS(FACILITIES).
        SEASONAL VARIATIONS. THUNDERSTORMS. VISIBILITY. WEATHER
        FORECASTING, WIND, DECISION MAKING,
DC (24) (11)
ID (25) Storm havens. La Spezia (italy).
IC (26)
AB (27)
        This handbook for the port of La Spezia, one in a
        series of severe weather duides for Mediterranean Sea
        ports, provides decision-making quidance for ship
        captains whose vessels are threatened by actual or
        forecast strong winds, high seas, restricted visibility
        or thunderstorms in the port vicinity. Causes and
        effects of such hazardous conditions are discussed.
        Precautionary or evasive actions are suggested for
        various vessel situations. The handbook is organized in
        four sections for ready reference: general quidance on
        nandbook content and use: a quick-look captain's
        summary: a more detailed review of deneral information
        on environmental conditions: and an appendix that
        provides oceanographic information. Keywords: Italian
        port facilities; Marine meteorology. Storm navens.
        Ligurian Sea. Ocean waves. Meteorological phenomena.
        Seāsonal variations. (edc)
AC (28) (U)
OL
   (33) 01
SE (34) F
   (35) 417277
```

Naval Environmental Prediction Research Facility

Monterey, CA 93943-5006

Contractor Report CR 88-09 May 1988



SEVERE WEATHER GUIDE: MEDITERRANEAN PORTS.

21. LA SPEZIA.

QUALIFIED REQUESTORS MAY OBTAIN ADDITIONAL COPIES FROM THE DEFENSE TECHNICAL INFORMATION CENTER.
ALL OTHERS SHOULD APPLY TO THE NATIONAL TECHNICAL INFORMATION SERVICE.

CONTENTS

Pre	eface	d	V
1.	GENE	RAL GUIDANCE	1-1
	1.1	Design	$1-1 \\ 1-1$
	1.2	Contents of Specific Harbor Studies	1-3
2.	Cap	otain's Summary	2-1
3.	Gene	ral Information	3-1
	3.1	Geographic Location	3-1
	3.2	Qualitative Evaluation of the Harbor as a Haven	3-5
	3.3	Currents and Tides	3-5
	3.4	Visibility	3-6
	3.5	Hazardous Conditions	3-6
	3.6	Harbour Protection	3-10
	3.7	Protective/Mitigating Measures	3-11
	3.8	Local Indicators of Hazardous Weather Conditions . 3.8.1 Genoa Lows, the Primary Cause of	
		Southeasterly Winds at La Specia	
		3.8.2 Scirocco	
	7 0		
	3.7	Summary of Problems, Actions, and Indicators	. 3-13
Re	feren	ces	. 3-25
Δη	nendi	x A General Purpose Oceanographic Information	Δ-1

FOREWORD

This handbook on Mediterranean Ports was developed as part of an ongoing effort at the Naval Environmental Prediction Research Facility to create products for direct application to Fleet operations. The research was conducted in response to Commander Naval Oceanography Command (COMNAVOCEANCOM) requirements validated by the Chief of Naval Operations (OP-096).

As mentioned in the preface, the Mediterranean region is unique in that several areas exist where local winds can cause dangerous operating conditions. This handbook will provide the ship's captain with assistance in making decisions regarding the disposition of his ship when heavy winds and seas are encountered or forecast at various port locations.

Readers are urged to submit comments, suggestions for changes, deletions and/or additions to Naval Oceanography Command Center (NAVOCEANCOMCEN), Rota with a copy to the oceanographer, COMSIXTHFLT. They will then be passed on to the Naval Environmental Prediction Research Facility for review and incorporation as appropriate. This document will be a dynamic one, changing and improving as more and better information is obtained.

W. L. SHUTT Commander, U.S. Navy

PORT INDEX

The following is a tentative prioritized list of Mediterranean Ports to be evaluated during the five-year period 1988-92, with ports grouped by expected year of the port study's publication. This list is subject to change as dictated by circumstances and periodic review.

1	GAETA, ITALY		BENIDORM, SPAIN
	NAPLES, ITALY		
3	CATANIA, ITALY		
4	AUGUSTA BAY, ITALY		PORT SAID, EGYPT
5	CAGLIARI, ITALY		ALEXANDRIA, EGYPT
6	LA MADDALENA, ITALY		ALGIERS, ALGERIA
7	MARSEILLE, FRANCE		TUNIS, TUNISIA
8	TOULON, FRANCE		GULF HAMMAMET, TUNISIA
9	VILLEFRANCHE, FRANCE		GULF OF GABES, TUNISIA
10	MALAGA, SPAIN		SOUDA BAY, CRETE
11	NICE, FRANCE		
12	CANNES, FRANCE	1991	PORT
13	MONACO		
14	ASHDOD, ISRAEL		PIRAEUS, GREECE
15	HAIFA, ISRAEL		KALAMATA, GREECE
16			THESSALONIKI, GREECE
17			CORFU, GREECE
18			KITHIRA, GREECE
19			VALETTA, MALTA
20	•		LARNACA, CYPRUS
21	· · · · · · · · · · · · · · · · · · ·		
22	VENICE, ITALY	1992	PORT
23	TRIESTE, ITALY		~~~~~~~~~~~~~~~~~
24			ANTALYA, TURKEY
25			ISKENDERUN, TURKEY
	SAN REMO, ITALY		IZMIR, TURKEY
	GENOA, ITALY		ISTANBUL, TURKEY
			GOLCUK, TURKEY
989 	PORT	_	GULF OF SOLLUM
	SPLIT, YUGOSLAVIA		
	DUBROVNIK, YUGOSLAVIA		
	TARANTO, ITALY		
	PALERMO, ITALY		

MESSINA, ITALY TAORMINA, ITALY PORTO TORRES, ITALY

PREFACE

Environmental phenomena such as strong winds, high waves, restrictions to visibility and thunderstorms can be hazardous to critical Fleet operations. The cause and effect of several of these phenomena are unique to the Mediterranean region and some prior knowledge of their characteristics would be helpful to ship's captains. The intent of this publication is to provide guidance to the captains for assistance in decision making.

The Mediterranean Sea region is an area where complicated topographical features influence weather patterns. Katabatic winds will flow through restricted mountain gaps or valleys and, as a result of the venturi effect, strengthen to storm intensity in a short period of time. As these winds exit and flow over port regions and coastal areas, anchored ships with large 'sail areas' may be blown aground. Also, hazardous sea state conditions are created, posing a danger for small boats ferrying personnel to and from port. At the same time, adjacent areas may be relatively calm. A glance at current weather charts may not always reveal the causes for these local effects which vary drastically from point to point.

Because of the irregular coast line and numerous islands in the Mediterranean, swell can be refracted around such barriers and come from directions which vary greatly with the wind. Anchored ships may experience winds and seas from one direction and swell from a different direction. These conditions can be extremely hazardous for tendered vessels. Moderate to heavy swell may also propagate outward in advance of a storm resulting in uncomfortable and sometimes dangerous conditions, especially during tending, refueling and boating operations.

This handbook addresses the various weather conditions, their local cause and effect and suggests some evasive action to be taken if necessary. Most of the major ports in the Mediterranean will be covered in the handbook. A priority list, established by the Sixth Fleet, exists for the port studies conducted and this list will be followed as closely as possible in terms of scheduling publications.

RECORD OF CHANGES

CHANGE NUMBER	DATE OF CHANGE	DATE ENTERED	PAGE NUMBER	ENTERED BY
			1	

1. GENERAL GUIDANCE

1.1 DESIGN

This handbook is designed to provide ship captains with a ready reference on hazardous weather and wave conditions in selected Mediterranean harbors. Section 2, the captain's summary, is an abbreviated version of section 3, the general information section intended for staff planners and meteorologists. Once section 3 has been read, it is not necessary to read section 2.

1.1.1 Objectives

The basic objective is to provide ship captains with a concise reference of hazards to ship activities that are caused by environmental conditions in various Mediterranean harbors, and to offer suggestions for precautionary and/or evasive actions. A secondary objective is to provide adequate background information on such hazards so that operational forecasters, or other interested parties, can quickly gain the local knowledge that is necessary to ensure high quality forecasts.

1.1.2 Approach

Information on harbor conditions and hazards was accumulated in the following manner:

- A. A literature search for reference material was performed.
- B. Cruise reports were reviewed.
- C. Navy personnel with current or previous area experience were interviewed.
- D. A preliminary report was developed which included questions on various local conditions in specific harbors.

- E. Port/harbor visits were made by NEPRF personnel; considerable information was obtained through interviews with local pilots, tug masters, etc; and local reference material was obtained.
- F. The cumulative information was reviewed, combined, and condensed for harbor studies.

1.1.3 Organization

The Handbook contains two sections for each harbor. The first section summarizes harbor conditions and is intended for use as a quick reference by ship captains, navigators, inport/at sea OOD's, and other interested personnel. This section contains:

- A. a brief narrative summary of environmental hazards.
- B. a table display of vessel location/situation, potential environmental hazard, effect-precautionary/evasion actions, and advance indicators of potential environmental hazards.
- C. local wind wave conditions, and
- D. tables depicting the wave conditions resulting from propagation of deep water swell into the harbor.

The swell propagation information includes percent occurrence, average duration, and the period of maximum wave energy within height ranges of greater than 3.3 feet and greater than 6.6 feet. The details on the generation of sea and swell information are provided in Appendix A.

The second section contains additional details and background information on seasonal hazardous conditions. This section is directed to personnel who have a need for additional insights on environmental hazards and related weather events.

1.2. CONTENTS OF SPECIFIC HARBOR STUDIES

This handbook specifically addresses potential wind and wave related hazards to ships operating in various Mediterranean ports utilized by the U.S. Navy. It does not contain general purpose climatology and/or comprehensive forecast rules for weather conditions of a more benign nature.

The contents are intended for use in both previsit planning and in situ problem solving by either mariners or environmentalists. Potential hazards related to both weather and waves are addressed. The oceanographic information includes some rather unique information relating to deep water swell propagating into harbor shallow water areas.

Emphasis is placed on the hazards related to wind, wind waves, and the propagation of deep water swell into the harbor areas. Various vessel locations/situations are considered, including moored, nesting, anchored, arriving/departing, and small boat operations. The potential problems and suggested precautionary/evasive actions for various combinations of environmental threats and vessel location/situation are provided. Local indicators of environmental hazards and possible evasion techniques are summarized for various scenarios.

CAUTIONARY NOTE: In September 1985 Hurricane Gloria raked the Norfolk, VA area while several US Navy ships were anchored on the muddy bottom of Chesapeake Bay. One important fact was revealed during this incident: Most all ships frigate size and larger dragged anchor, some more than others, in winds of over 50 knots. As winds and waves increased, ships 'fell into' the wave troughs, BROADSIDE TO THE WIND and become difficult or impossible to control.

This was a rare instance in which several ships of recent design were exposed to the same storm and much effort was put into the documentation of lessons learned. Chief among these was the suggestion to evade at sea rather than remain anchored at port whenever winds of such intensity were forecast.

2. CAPTAIN'S SUMMARY

The Port of La Spezia is located on the northeast shoreline of the Ligurian Sea at 44°05'N 09°51'E, near the eastern limit of the Gulf of Genoa (Figure 2-1).




Figure 2-1. Ligurian Sea.

The spacious port is situated at the head of Golfo di La Spezia (Gulf of La Spezia), and is protected from most hazardous weather conditions. Constructed around a natural inlet with approximate dimensions of 10,000 ft (3048 m) wide by 15,000 ft (4,572 m) long, the port can easily accommodate over 40 medium— to deep—draft vessels (FICEURLANT, 1987). See Figure 2-2.




Figure 2-2. Region of the Ports of Livorno and La Spezia.

The Port of La Spezia is divided into an outer harbor and inner harbor by two breakwaters. The outer breakwater, Diga Foranea, is 7,218 ft (2,200 m) long, and forms the southern limit of the outer harbor (Figure 2-3). Two entrances to the harbor exist, with one at each end of Diga Foranea. The western entrance, with a depth of 44 ft (13.5 m) is most often used.

The inner harbor, Darsena Duca Degli Abruzzi (Porto Militare—the Italian Navy Base) is formed by the second breakwater system in the northwestern part of the Port. Small surface ships (frigate size or smaller) and submarines can "Med-moor" in the inner harbor. Entrance depth to the inner harbor is 36 ft (11 m) but is subject to silting. (In 1987, a U.S. submarine scraped bottom while entering the basin.) Fleet landing is in the Duca Degli Abruzzi at the Bachino di Revello.

Several anchorages exist at La Spezia. One is located inside the breakwater approximately 2,000 yd (1,829 m) northwest of Diga Foranea in a depth of 34.5 ft (10.5 m). Holding is rated good on a mud bottom. Two unprotected (exposed to open sea conditions) anchorages are located outside the breakwater. One is 2 n mi south (depth 86 ft) and the other is 0.6 n mi east (depth 66 ft) of Tino Island. The anchorages provide good holding on mud bottoms.




Figure 2-3. Port of La Spezia.

According to the <u>Port Directory</u> (FICEURLANT, 1987), most currents in the harbor are wind generated, but a weak counterclockwise current usually prevails. Tides within the harbor have a maximum variation of about 20 inches (0.5 m).

Specific hazardous conditions, vessel situations, and suggested precautionary/evasive action scenarios are summarized in Table 2-1.

Table 2-1. Summary of hazardous environmental conditions for the Port of La Spezia, Italy.

HAZARDOUS CONDITIONS	INDICATORS OF POTENTIAL HAZARD	VESSEL LOCATION/ SITUATION AFFECTED	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS
. N'ly winds/waves † Mind locally called Tramontana. † Maximum velocities normally force 6 to 7 (22 to 33 kt). † Mind raises 3 ft sea in outer harbor. † Waves reflect off N side of outer breakwater, creating a chop in nearby waters.	Advance warning * Caused by building high pressure over N Italy as low pressure center moves SE of La Spezia. Duration * To be expected as long as strong high pressure dominates N Italy. * Will likely weaken quickly when signs of Genoa cyclogenesis appear.	(1) Moored - Italian Navy Base. (2) Anchored - outer harbor.	(a) Minimal effect - SHIPS HAVE NEVER BEEN MOVED DUE TO BAD MEATHER. † Additional mooring lines may be required in a strong event. (b) Be aware of wind chill factor. (a) Minimal effect - SHIPS HAVE NEVER BEEN MOVED INSIDE THE BREAKWATER DUE TO BAD MEATHER. * Ships anchored near the outer breakwater may experience unconfortable motion due to reflected waves.
		(3) <u>Anchored – outside breakwater</u> .	(b) Be aware of wind chill factor. (a) Minimal effect - due to limited fetch length. * Small boat runs to/from anchorage may be affected.
		(4) <u>Arriving/departing</u> .	(b) Be aware of wind chill factor. (a) Minimal impact. * Vessels inbound to moorings at the Italian Navy Base may need extra mooring lines. * Vessels anchoring near outer breakwater should be aware of vessel motions induced by reflected waves.
		(5) <u>Small boats</u> .	(b) Be aware of wind chill factor. (a) Small boat operations should be largely unaffected in the inner and outer harbors. * Small boat operators should be aware of hazards posed by reflected waves near the N side of the outer breakwater. * Runs to/from outer anchorages may be curtailed. * Local water taxis will not go outside the breakwater in winds of force 6 (22-27 kt) or greater.
SE'ly winds/waves • Wind locally called Scirocco, but	Advance warning * Commonly associated with low pressure systems	(1) <u>Moored - Italian Navy Base</u> .	(b) Be aware of wind chill factor. (a) Minimal effect - SHIPS HAVE NEVER BEEN MOVED DUE TO BAD WEATHER. † Additional mooring lines may be required in a strong event.
true Scirocco events occur only infrequently. * Maximum velocities normally force 7 to 8 (28 to 40 kt). * May occur 3-4 times per year. * If wind direction is SSE or S, waves may be raised in the outer anchorages.	occur only normally force t). s per year. SSE or S, waves acour give on the Gulf of Genoa. acour almost sizultaneously with the onset of a Mistral in the Gulf of Sper year. Lion, and invariably form when conditions	(2) <u>Anchored - outside breakwater</u> .	(a) Minimal effect, unless wind is primarily from SSE or S. * SE wind would have insufficient fetch to generate hazardous waves. * SSE or S wind would have a trajectory which would cause significant waves to generate and reach the anchorages. * Evasion at sea is recommended if strong winds are forecast. * No nearby ports offer better protection.
	through France reaches a lee trough which is often present in the Bulf of Benoa. * May be associated with a slowly developing Scirocco event. * Clouds forming on SE slope of Mt. Parodi may indicate the beginning of a Scirocco. Duration	(3) <u>Arriving/departing</u> .	 (a) Minimal impact on inbound/outbound vessels. * Vessels inbound to moorings at the Italian Navy Base may need extra lines. * Anchoring outside the breakwater may not be feasible due to high waves if the wind is from SSE or S (Vice SE). * Small boat runs to/from the outer anchorage may be curtailed in a strong event. Local water taxis will not go outside the breakwater in winds of force 6 (22-27 kt) or greater.
	* A strong event may last for up to 3 days. * Commonly lasts a day or less. Associated weather * Convective activity associated with a Benoa low has a periodicity of 18 hr, starting with cold frontal passage. * Most intense activity occurs at 36 hr intervals. * Dust and/or 'red rain' often accompany Scirocco winds.	(4) <u>Small boats</u> .	(a) Minimal impact on operations inside the breakwater. # BOATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER DUE TO BAD WEATHER. (b) Boat to/from the outer anchorage may be affected. # Waves may preclude safe boat operation in a strong event, or if the wind direction is from SSE or S (vice SE). # Local water taxis will not go outside the breakwater in winds of force 6 (22-27 kt) or greater.

Table 2-1. (Continued)

HAZARDOUS CONDITIONS	INDICATORS OF POTENTIAL HAZARD	VESSEL LOCATION/ SITUATION AFFECTED	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS
3. SN'ly winds/waves * Winds with strong N component reach the harbor as SN after passing through valleys in peninsula N of the Port. * Normally limited to force 5 (17-21 kt). * Any wind from SSN to NNN has potential to generate waves that would adversely impact the outer anchorages.	Advance warning * Cyclogenesis or low pressure center moving over N Italy or Bulf of Venice. * Cold/occluded frontal passage. Duration * Likely of short duration as winds will tend to veer through N to NN or N as pressure builds behind low or front.	(1) Anchored - outside breakwater. (2) Arriving/departing.	(a) The anchorage may be unusable in heavy weather. * Evasion at sea is recommended if heavy weather is forecast. * No nearby ports offer better protection. (a) Minimal impact on inbound/outbound vessels. * Vessels will be protected from open-sea conditions when in the lee of Palaeria Island, but the protection is lost when south of the island. * Tino Island offers only limited protection close alee. * Outer anchorages are exposed and should be discounted as alternatives if heavy weather is forecast.
		(3) <u>Small boats</u> .	(a) Operations inside the breakwater are only minimally affected. * BOATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER DUE TO BAD WEATHER. (b) Runs to/from the outer anchorages may be curtailed. * Anchorages are exposed to open-sea conditions. * Local water taxis will not go outside the breakwater in winds of force 6 (22-27 kt) or greater.

SEASONAL SUMMARY OF LA SPEZIA HAZARDOUS WEATHER CONDITIONS

WINTER (November thru February)

- * NW'ly winds: called Tramontana, may reach force 6-7 (22-33 kt), result of building high pressure over Italy as low pressure moves SE from Gulf of Genoa.
- * SE'ly winds: called Scirocco caused by low pressure developing in Gulf of Genoa. Speeds up to force 7-8 (28-40 kt) expected 3-4 times/year. True Scirocco accompanied by North African dust particles ("red rain").
- * W'ly winds: funneled thru two valleys on west side of port, experienced at port SW'ly (up to force 5 (17-21 kt).

SPRING (March thru May)

- * High winds (99 kt April and 63 kt May) from unspecified direction have been recorded.
- * Tramontana (NW'ly winds) and Scirocco (SE'ly winds) possible. Strongest/most frequent earlier in season.
- * Sea breeze possible (afternoon).

SUMMER (June thru September)

- * Sea breeze (light to moderate).
- * Scattered thunderstorms.
- * Small waterspouts and small tornadoes have been reported.

AUTUMN (October)

- * Transition month.
- * Tramontana (NW'ly winds) and Scirocco (SE'ly winds) increasing in strength and frequency.

NOTE: For more detailed information on hazardous weather conditions see previous Summary Table in this section and Hazardous Weather Summary in Section 3.

REFERENCES

FICEURLANT, 1987: <u>Port Directory for La Spezia (1984)</u>, <u>Italy</u>. Fleet Intelligence Center Europe and Atlantic, Norfolk, VA.

3. GENERAL INFORMATION

This section is intended for Fleet meteorologists/oceanographers and staff planners. Paragraph 3.5 provides a general discussion of hazards, and Table 3-2 provides a summary of vessel locations/situations, potential hazards, effect - precautionary/evasive actions, and advance indicators and other information about potential hazards by season.

3.1 Geographic Location

The Port of La Spezia is located on the northeast shoreline of the Ligurian Sea at 44°05'N 09°51'E, near the eastern limit of the Gulf of Genoa (Figure 3-1).




Figure 3-1. Ligurian Sea.

The spacious port is situated at the head of Golfo di La Spezia (Gulf of La Spezia), and is protected from most hazardous weather conditions. It is constructed around a natural inlet which has approximate dimensions of 10,000 ft (3,048 m) wide by 15,000 ft (4,572 m) long (Figure 3-2). It can easily accommodate over 40 medium— to deep—draft vessels (FICEURLANT, 1987).




Figure 3-2. Region of the Ports of Livorno and La Spezia.

The Port of La Spezia is divided into an outer harbor and inner harbor by two breakwater systems. outer breakwater, Diga Foranea, the surface of which is awash (Hydrographer of the Navy, 1965), is 7,218 ft (2,200 m) long, and forms the southern limit of the outer harbor (Figure 3-3). Two entrances to the harbor exist, with one at each end of Diga Foranea. The western entrance has a depth of 44 ft (13.5 m) and is most often The eastern entrance has a depth of (11.5 m), but is often congested with drydock activity and moored tankers. The inner harbor, Darsena Duca Degli Abruzzi (Porto Militare--the Italian Navy Base) is formed by the second breakwater system in the northwestern part the Port. Small surface ships (frigate smaller) and submarines can "Med-moor" in the inner harbor. Entrance depth to the inner harbor is 36 ft (11 m), but is subject to silting. (In 1987 a U.S. submarine scraped bottom while entering the basin.) Fleet landing is in the Duca Degli Abaruzzi at the Bachino di Revello.

Other facilities located inside the outer breakwater include commercial facilities along the north and east perimeters, and Seno Della Grazie, a cove located northwest of the western entrance to the outer harbor. Seno Della Grazie is used as a berthing place for small passenger liners (FICEURLANT, 1987).

Several anchorages exist at La Spezia. One protected anchorage is located in the outer harbor approximately 2,000 yd (1,829 m) northwest of Diga Foranea in a depth of 34.5 ft (10.5 m). Holding is rated good on a mud bottom. Two open sea (unprotected) anchorages are located outside the breakwater. One is located 2 n mi south (depth 86 ft) and 0.6 n mi east (depth 66 ft) of Tino Island. The anchorages have good holding qualities on mud bottoms.




Figure 3-3. Port of La Spezia.

3.2 Qualitative Evaluation of Harbor as a Haven

The Port of La Spezia is effectively protected on three sides by steeply rising terrain. Elevations close westward of the port commonly exceed 1,500 ft (457 m), with Mt. Verrugoli reaching 2,444 ft (745 m). East of the port, elevations are somewhat lower, but some peaks exceed 1,000 ft (305 m) and one peak, Mt. Rocchetta, reaches 1,352 ft (412 m). The terrain protects the port from many high wind situations. but valleys which penetrate the mountainous areas allow some wind events to reach the harbor. Anchorages in the harbor and berths in the Italian Naval Base are relatively safe. According to local authorities, SHIPS HAVE NEVER HAD TO BE MOVED DUE TO BAD WEATHER, AND BOATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER. Anchorages outside the breakwater are exposed, however, and evasion at sea is recommended in heavy weather. Water taxis will not go to the outside anchorages when winds are force 6 (22-27 kt) or greater.

The inner and outer harbors are well protected from significant wave action with the exception that strong northerly winds raise a sea which reflects off the north face of Diga Foranea and causes uncomfortable, though not dangerous, motion to ships anchored nearby.

3.3 Currents and Tides

According to the <u>Port Directory</u> (FICEURLANT, 1987), most currents are wind generated. Local authorities state that a weak counter-clockwise current exists in the harbor. Similarly, <u>Mediterranean Pilot</u> (Hydrographer of the Navy, 1965) states that normally the same current exists in the whole of Golfo di La Spezia, but the current is strongly influenced by the prevailing winds.

Tides within the harbor are minimal, with a maximum range of about 20 inches (0.5 m).

3.4 Visibility

Visibility is normally good all year, but salt haze reduces visibility to 5 or 6 n mi during summer. A review of visibility statistics for the period 1951-1979 reveals that visibility restricted below 2.5 mi has the highest frequency of occurrence during the months of October through May.

3.5 Hazardous Conditions

La Spezia is located adjacent to one of the most active areas of cyclogenesis in the world, the Gulf of Genoa. Consequently, it is subject to frequent periods of unstable, inclement weather, mainly during the autumn, winter and spring seasons.

A seasonal summary of various known environmental hazards that may be encountered in the Port of La Spezia follows.

A. Winter (November through February)

Al though northerly Tramontana winds experienced during all seasons of the year, they are most frequent and stronger during winter. Maximum velocities are normally force 6 to 7 (22-33 kt). The winds have no effect at dockside due to the excellent protection afforded by the surrounding mountainous terrain. north of and near to Diga Foranea, breakwater which defines the southern limit of the outer experience uncomfortable, though normally innocuous, vessel motion as northerly wind waves reflect off the north side of the breakwater. Strong Tramontana winds are the result of building high pressure over Italy as a 1 ow pressure center moves southeastward from the Gulf of Genoa.

Southeasterly winds are also experienced year round, abut the highest speeds are observed during winter. Although locally referred to as Scirocco, most are not of true Scirocco origin, but are instead caused by low pressure systems developing in or moving into the

Gulf of Genoa. Speeds of force 7 to 8 (28-40 kt) can be expected 3 or 4 times per year, with late winter episodes likely. Although such occurrences are relatively strong, wave heights inside the breakwater are limited to near 3 ft (about 1 m). The infrequent strong events may last up to 3 days, but the usual duration is a day or less.

True Scirocco winds are infrequently observed at La Spezia. When they occur, they are normally accompanied by dust and/or "red rain" as wind-borne particles from North Africa reach the area. Wind speeds are not normally strong.

Westerly wind regimes are experienced in the port area as southwesterly. Speeds up to force 5 (17-21 kt) occur as the winds funnel through two valleys that penetrate the steep terrain west of the Port. Effect on the Port is minimal.

Precipitation is common during winter. November has the greatest monthly accumulation of the year, about 4 inches. About 6 days in each month of the season, amounts from a trace (.01 in) to 0.4 in (9.9 mm) are recorded. Amounts of 0.4 inches (10 mm) or more are normally recorded on approximately 3 days of each month. While not frequently observed, snow is possible during the months of December, January, and February.

Temperatures during the coldest month of the season, January, range from a median low of 41°F (5°C) to a median high of 51°F (11°C). The record low temperature recorded during a 29 year period of record ending in 1979 was 20.8°F (-6.2°C). Because of the relatively cold temperatures that are common at La Spezia, the effect of wind chill (temperature combined with wind speed) should be considered for evolutions requiring personnel to work in exposed locations. Table 3-1 can be used to determine the wind chill factor for various temperature and wind speed combinations.

Table 3-1. Wind Chill. The cooling power of the wind expressed as "Equivalent Chill Temperature" (adapted from Kotsch. 1983).

Wind S		_					•	essec	d as	
Knots	MPH		"Equivalent Chill Temperature" Temperature (°F)							
Calm	Calm	40	35	30	25	20	15	10	5	0
			Equi	vale	ent (Chill	. Ten	npera	ature	•
3-6	5	35	30	25	20	15	10	5	0	-5
7-10	10	30	20	15	10	5	0	-10	-15	-20
11-15	15	25	15	10	0	-5	-10	-20	-25	-30
16-19	20	20	10	5	0	-10	-15	-25	-30	-35
20-23	25	15	10	0	-5	-15	-20	-30	-35	-45
24-28	30	10	5	0	-10	-20	-25	-30	-40	-50
29-32	35	10	5	-5	-10	-20	-30	-35	-40	-50
33-36	40	10	0	-5	-15	-20	-30	-35	-45	-55

B. Spring (March through May)

The spring season is characterized by periods of stormy, winter-type weather which alternates with a number of false starts of relatively settled summer-type weather (Brody and Nestor, 1980). The more unsettled periods occur during March and April, when low pressure systems develop in or pass through the Gulf of Genoa. After April, the transition to settled weather proceeds more smoothly. Although strong winds are not the rule at Spezia, a potential for damaging winds exists. possibility is evidenced by the fact that the climatology record from Palmaria Island (Figure 3-3) speeds of 99 kt have been recorded during April. 62 kt during May. The direction was not specified in either instance, but the existence of such a wind in the past indicates the possibility of a similar event in the future cannot be discounted.

Northerly Tramontana winds are possible throughout the season, but are strongest early in the period because the high pressure systems over northern Italy become less strong as the season progresses.

Southeasterly winds are also experienced throughout the season, and like the Tramontana, are strongest early in the season when Genoa lows are the strongest.

Scirocco events are possible early in the season and can bring dust and/or "red rain" to La Spezia.

Prevailing night and early morning winds are northerly, but southwesterly winds are the rule by midafternoon as a sea breeze regime becomes evident on warm days early in the season.

Precipitation amounts remain near the wintertime levels through March, but decrease thereafter.

Temperatures warm throughout the season as summer approaches. By May, the mean daily maximum temperature has risen to 67°F (19°C), while the minimum has risen to 55°F (13°C). Wind chill factor is a consideration until late April. See Table 3-1.

C. Summer (June through September)

Summer is a period of relatively warm, settled conditions at La Spezia. Strong winds are uncommon, but velocities of 60 kt have been recorded during August, and 55 kt during September.

Night and early morning winds are normally from the north, with southwesterly winds common during the afternoon. Warmest temperatures are observed during July and August, with a mean daily maximum of 79°F (26°C). Precipitation is at its yearly minimum in July, when less than 1 inch of rain accumulates during an average year. The monthly total increases to almost 3 inches by September.

La Spezia experiences scattered thunderstorms during the summer. Small waterspouts and tornadoes have also been reported.

D. Autumn (October)

Autumn usually lasts for the single month of October and is characterized by an abrupt change to winter-type weather (Brody and Nestor, 1980).

The possibility of stronger northerly Tramontana wind increases as influxes of cold air begin to invade Europe in advance of the winter season. Similarly, the threat of stronger southeasterly winds increases as cyclonic activity in the Gulf of Genoa starts to become more common.

Temperatures decrease from those of summer, with a mean maximum of about 67°F (19°C) and mean minimum of 55°F (13°C) normal for October. Precipitation amounts increase from those of September, with an average accumulation of 3.76 inches occurring on about 7.5 days during the month. Thunderstorms occur on an average of 3 days during October, the greatest frequency of the year.

3.6 Harbor Protection

The Port of La Spezia is afforded excellent protection from most wind and wave conditions, but is vulnerable to certain specific events. Also, as detailed below, there are differences in protection enjoyed by vessels inside the breakwaters and those anchored in open sea conditions.

3.6.1 Wind and Weather

The inner and outer harbors are bordered on 3 sides (west, north, and east) by steeply rising terrain which serves to block or reduce the speed of many wind regimes. Relatively strong winds reach the port, however, from the southeast, southwest, and north. Anchorages in the harbor and berths in the Italian Naval Base (Darsena Duca Degli Abruzzi) are relatively safe; SHIPS HAVE NEVER HAD TO BE MOVED DUE TO BAD WEATHER AND BOATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER. However, water taxis will not go to the outside anchorages when winds are force 6 (22-27 kt) or greater.

The harbor affords no effective protection from thunderstorms nor the infrequently seen small waterspouts and tornadoes.

3.6.2 Waves

Inside the breakwater, wave action is limited to those which are raised by the local wind. Since the maximum straight line distance across the harbor inside the breakwater (Diga Foranea) is only 2.5 n mi, wave generation is fetch limited. Southerly winds of force 7 to 8 (28-40 kt) only raise a 3 ft (about 1 m) sea inside the breakwater, so operations are only minimally affected. The same situation applies to northerly winds, but with a different result. According to local authorities, northerly waves reflect off the north face of Diga Foranea and cause "uncomfortable, though not usually damaging", vessel motion to ships anchored near the breakwater.

Vessels anchored in the two outside anchorages are exposed to open sea conditions. Consequently, any wave condition which emanates from south through northwest has the potential to reach the anchorages.

3.7 Protective/Mitigating Measures

3.7.1 Sortie/Remain in Port

Vessels at the Italian Naval Base should be able to remain in Port. SORTIE HAS NEVER BEEN NECESSARY DUE TO BAD WEATHER IN THE PORT OF LA SPEZIA.

3.7.2 Moving to a New Anchorage

Vessels anchored in the harbor should be able to stay; SHIPS HAVE NEVER HAD TO BE MOVED DUE TO BAD WEATHER. But because they are exposed to open-sea conditions, ships in the two anchorages outside the breakwater, 0.6 n mi east and 2 n mi south of Tino Island, should evade at sea if heavy weather is forecast. There are no good alternate anchorages near La Spezia in which to seek haven.

3.8 Local Indicators of Hazardous Weather Conditions

The Port of La Spezia affords excellent shelter from most hazardous weather conditions, but it is prudent to be aware of potential hazards and have insight as to

when specific conditions may develop. The following guidelines have been extracted from various sources, including on-site interviews with local authorities. They are intended to provide additional insight to the Fleet meteorologists, and enable them to recognize events that indicate changes in weather conditions. Unless otherwise indicated the guidelines have been adapted from Regional Forecasting Aids for the Mediterranean Basin, (Brody and Nestor, 1980).

3.8.1 Genoa Lows, the Primary Cause of Southeasterly Winds at La Spezia.

A. Cyclogenesis

- 1. A lee trough often is present in the Gulf of Genoa when a cold or occluded front is moving into western France. This lee trough remains stationary until the arrival of the front, at which time significant cyclogenesis occurs.
- 2. A good indication of rapid development of a Genoa low is the appearance of cold air from the northeast in the Po Valley of northern Italy.
- 3. If Genoa cyclogenesis is predicted, the following rules can be used to decide whether it will occur in the Gulf of Genoa or to the east in the Gulf of Venice:
- a. If large amounts of cold air penetrate the Po Valley from the northeast, cyclogenesis can be expected in the Gulf of Genoa. This cyclone generally will move southeastward along the west coast of Italy.
- b. If little cold air penetrates the Po Valley from the northeast while a strong push is observed in the Gulf of Lion, cyclogenesis will probably take place in the Gulf of Venice. This cyclone occasionally may move southeast through the Adriatic Sea.
- 4. Genoa lows occur almost simultaneously with the onset of the Mistral in the Gulf of

Lion, and invariably form when conditions are right for the Mistral to occur.

5. Complex low pressure systems with multiple centers at the surface are a common event in the western Mediterranean Basin. One center usually can be found in the gulf of Genoa, while another is found over North Africa; a weak pressure gradient exists between the two systems. Which of these lows will develop depends greatly on the movement of an upper-level (500 mb) short wave trough. If the trough moves to the North African coast, for example, the low center in that region will develop rapidly, increasing the pressure gradient and causing easterly gales over the southern Tyrrhenian Sea.

B. Associated wind and weather

- 1. Weak to moderate Genoa cyclogenesis causes important variations in the weather along the west coast of Italy. When analyzing these cases, the resolution of the 500 mb analysis should be fine enough to support tracking of the weak short wave troughs associated with increased shower activity.
- 2. Convective activity associated with a Genoa low has a periodicity of about 18 hr, starting with the initial cold frontal passage. The periodicity is most pronounced with a stationary low. The most intense convective activity occurs at 36 hr intervals.
- 3. Strong westerly winds associated with Mistral conditions rarely reach the west coast of Italy. Winds usually will not reach gale force until after the associated Genoa low moves off to the southeast. Under these conditions gale force northeasterly winds occur along the west coast of Italy.
- 4. Strong northerly winds can be expected in the Gulf of Genoa within 6-8 hr if (1) the 1034 mb isobar is present along the crest of the Alps north of the Gulf of Genoa and (2) increasing northerly winds are observed at Milan (16080).

C. Miscellaneous

 A residual low pressure trough generally remains over the Gulf of Genoa even after the primary low has moved well out of the region. The trough can remain for several days.

2. Centers of Genoa lows can be poorly organized; strong pressure gradients, associated with a lee trough south of the Alps, frequently are found far from the low's geographic center.

3.8.2 Scirocco

A. During Scirocco periods over the Mediterranean, a low-level jet is likely to exist just below the top of the very marked temperature inversion common during the Scirocco. Wind speeds reaching 70-80 kt, with heavy turbulence associated with the strong vertical wind shear, have been observed in the jet.

B. Local indicators

The following guidelines were related by local authorities:

- 1. Scirocco winds usually will increase gradually, so ample advance warning is provided by the time taken to develop a full Scirocco situation.
- 2. Clouds forming on the southeast slope of Mount Parodi (northwest of La Spezia) may indicate the beginning of a Scirocco.

3.8.3 General

- A. Past experience indicates that the reliability of hourly reporting stations along the Italian coast is questionable, especially during the night.
- B. During periods of southerly surface flow in the central Mediterranean, convergence zones between southeasterly and southwesterly winds are frequently observed. These convergence zones result in heavier precipitation and lower visibilities. Fronts are not associated with this phenomenon initially, but may develop later.

- c. During the winter half of the year along the west coast of Italy, maximum occurrence of convective activity is in the early morning (0300-0800L) and minimum occurrence is in the late afternoon and early evening. In the mountains to the east, however, this diurnal variation is reversed.
- D. Dry, moderate-to-strong (15-25 kt), north-to-east winds during the winter have produced steam fog along the Italian coast from Genoa to Pisa, out to 35 n mi offshore. Visibilities in this fog are reduced to 1-2 n mi although the dewpoint-temperature spread measured at an aircraft carrier's flight deck level may exceed 4°F.
- E. Local authorities state that clouds forming on the southeast slope of Mount Parodi (northwest of La Spezia) may indicate the beginning of a period of bad weather and rain is about to start.

3.9 Summary of Problems, Actions, and Indicators

Table 3-2 is intended to provide easy to use seasonal references for meteorologists on ships using the Port of La Spezia. Table 2-1 (section 2) summarizes Table 3-2 and is intended primarily for use by ship captains.

Table 3-2. Potential problem situations at Port of La Spezia, Italy - ALL SEASONS.

VESSEL LOCATION/SITUATION	POTENTIAL HAZARD	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS	ADVANCE INDICATORS AND OTHER INFORMATION ABOUT POTENTIAL HAZARD
1. Moored - Italian Navy Base. Strongest in Winter Occurs in Spring Summer and Autoan	a. N'ly winds - Locally called Tramontana. Caused by building high pressure over contern Italy as a low pressure center moves SE from the Gulf of Benoa. Maximum winds normally force 6 to 7 (22-33 kt).	a. Minimal effect. Additional mooring lines may be required in a strong event. SHIPS HAVE NEVER BEEN MUVED DUE TO BAD WEATHER. Be aware of wind chill factor.	a. N'ly winds are the result of building high pressures over N Italy and a SE'ly moving low pressure center.
Strongest in Winter - Occurs in Spring Summer and Autumn	b. SE'ly winds - Locally called Scirocco. Caused by low pressure systems developing in or moving into the Bulf of Genoa. Velocities to force 7 to 8 (28-40 kt) can be expected 3-4 times per year. Stronger events may last up to 3 days, but the usual duration is a day or less. True Scirocco events occur infrequently and are normally accompanied by dust and/or "red rain" as wind borne particles from North Africa reach the area.	b. Minimal effect. Additional mooring lines may be required in a strong event. SHIPS HAVE NEVER BEEN MOVED DUE TO BAD MEATHER.	b. SE'ly winds at La Spezia are most often caused by low pressure systems located in the Bulf of Benoa, but are infrequently caused by Scirocco events. (1) Senoa lows. The following is an abbreviated listing of various aspects of Benoa low behavior: (a) Cyclogenesis (b) Cyclogenesis 1. A fee trough often is present in the Bulf of Benoa when a cold or occlude front is moving into western France. The trough remains stationary until arrival of the front, at which time significant cyclogenesis occurs. 2. A good indication of rapid development of a Benoa low is the appearance of cold air from the NE in the Po Valley. Mistral in the Bulf of Linn, and invariably form when conditions are right for a Mistral in the Bulf of Linn, and invariably form when conditions are right for a Mistral to occur. (b) Associated weather: Convective activity associated with a Benoa low has a periodicity of about 18 hr, starting with the initial cold frontal passage, and is most pronounced with a stationary low. The most intense convective activity occurs at 36 hr intervals. (c) Miscellaneous Bulf of Benoa even after the primary low has moved well out of the region. The trough can remain several days. 2. Centers of Benoa lows can be poorly organized: Strong pressure gradients frequently are found far from the low's geographic center.
2. <u>Anchored - outer harbor</u> . Strongest in Winter Occurs in Spring Summer and Autuan	a. <u>K'ly winds/waves</u> - Locally called Tranontana. Caused by building high pressure over northern Italy as a low pressure center moves SE from the Bulf of Genoa. Maximum winds normally force 6 to 7 (22-33 kt). Raises a N'ly sea to 3 ft in the outer harbor which reflects off of the N side of the outer breakmater and creates a chop in the nearby maters.	a. Wind alone causes no significant problem, but reflected waves cause uncoafortable, but not usually damaging, vessel motions to vessels anchored near the breakwater. SHIPS HAVE NEVER BEEN MOVED DUE TO BAD MEATHER. Be aware of wind chill factor.	(2) Scirocco events (a) Clouds forming on the SE slope of Mt. Parodi (NW of La Spezia) may indicate the beginning of a Scirocco. (b) Scirocco winds usually will increase slowly, so ample advance warning is provided by the time taken to develop a full Scirocco situation. a. N'ly winds are the result of building high pressures over N Italy and a SE'ly moving low pressure center.

Table 3-2. (Continued)

VESSEL LOCATION/SITUATION	POTENTIAL HAZARD	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS	ADVANCE INDICATORS AND OTHER INFORMATION ABOUT POTENTIAL HAZARD
3. Anchored - outside breakwater. Strongest in Winter Occurs in Spring and Autumn Weakest in Summer	a. N'ly winds - Locally called Tramontana. Caused by building high pressure over northern Italy as a low pressure center moves SE from the built of Genoa. Haximum winds normally force 6 to 7 (22-33 kt).	a. Configuration of coastline minimizes the impact of wind and waves. Winds blowing S toward anchorages would have insufficient fetch length to raise a sea hazardous to anchored vessels. Saall boat runs to/from the inner harbor may be affected. Be aware of wind chill factor.	a. M'ly winds are the result of building high pressures over N Italy and a SE'ly moving low pressure center.
Strongest in Winter Occurs in Spring and Autumn Weakest in Summer	b. SE'ly winds - Locally called Scirocco. Caused by low pressure systems developing in or moving into the Gulf of Benoa. Velocities to force? to 8 128-40 kt) can be expected 3-4 times per year. Stronger events may last up to 3 days, but the usual duration is a day or less. True Scirocco events occur infrequently and are normally accompanied by dust and/or 'red rain' as wind borne particles from North Africa reach the area.	b. Configuration of coastline sinimizes the effect of any wind with a significant E component, but winds from SSE or S could raise maves that would impact the anchorages. Evasion at sea is recommended if heavy weather is forecast.	b. SE'ly winds at La Spezia are most often caused by low pressure systems located in the Gulf of Benoa, but are infrequently caused by Scirocco events. (1) Genoa low behavior. (a) Cyclogenesis 1. A lee trough often is present in the Gulf of Benoa when a cold or occluded front is moving into mestern France. The trough remains stationary until arrival of the front, at which time significant cyclogenesis occurs. 2. A good indication of rapid development of a Benoa low is the appearance of cold air from the NE in the Po Valley. 3. Benoa lows occur aloust simultaneously with the onset of a Mistral in the Bulf of Lion, and invariably form when conditions are right for a Mistral to occur. (b) Associated weather: Convective activity associated with a Genoa low has a periodicity of about 1B hr, starting with the initial cold frontal passage, and is aost pronounced with a stationary low. The most intense convective activity occurs at 36 hr intervals. (c) Miscellaneous 1. A residual low pressure trough generally remains over the Bulf of Benoa even after the primary low has moved well out of the region. The trough can remain several days. 2. Centers of Benoa lows can be poorly organized: Strong pressure gradients frequently are found far from the low's geographic center. (2) Scirocco events (a) Clouds forsing on the SE slope of Mt. Parodi (NW of La Spezia) may indicate the beginning of a Scirocco. (b) Scirocco winds usually will increase slowly, so ample advance warning is provided by the time taken to develop a full Scirocco situation.
Strongest in Winter Occurs in Spring Summer and Autumn	c. SW'ly winds/waves — Any relatively strong wind with a direction from SW through WNM has the potential to generate sea/swell that would adversely impact the anchorages. Any wind with a strong W component would reach the inner harbor as SW after passing through valleys in peninsula W of the Port. Normally limited to force 5 (17-21 kt) in harbor area. May be caused by low pressure N and/or Nc of La Spezia or as post frontal winds following passage of a cold/occluded front.	c. The anchorages are exposed to winds/waves from SSE through WNW. Evasion at sea is recommended if heavy weather is forecast.	c. Cyclogenesis or low pressure center moving over N Italy or Gulf of Venice may result in SM wind at La Spezia. SM to N winds may follow the passage of a cold or occluded front. If high pressure builds in behind a SE'ly moving low pressure center, the winds will likely continue to veer, becoming N as the low moves.

Table 3-2. (Continued)

VESSEL OCATION/SITUATION	POTENTIAL HAZARD	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS	ADVANCE INDICATORS AND OTHER INFORMATION ABOUT POTENTIAL HAZARD
4. Arriving/departing. Strongest in Winter Occurs in Spring and Autuan Weakest in Susmer	a, N'ly winds/waves - Locally called framontana. Caused by building high pressure over northern Italy as a low pressure center moves SE from the Bulf of Benoa. Maximum winds normally force 6 to 7 (22-33 kt). Raises a N'ly sea to 3 ft in the outer harbor which reflects off the N side of the outer breakwater and creates a chop in the nearby waters.	a. Inbound/outbound vessels should not encounter any significant problems while entering or leaving the Port. Ships inbound to the Italian Navy Base may need extra lines for mooring. Ships intending to anchor in the outer harbor near the breakwater should be aware of the bothersome chop created by reflected wind waves. Be aware of wind chill factor.	a. N'ly winds are the result of building high pressures over N Italy and a SE'ly moving low pressure center.
Strongest in Winter Occurs in Spring and Autumn Weakest in Summer	b. SE'ly winds - Locally called Scirocco. Caused by low pressure systems developing in or moving into the Gulf of Genoa. Velocities to force 7 to 8 (28-40 kt) can be expected 3-4 times per year. Stronger events may last up to 3 days, but the usual duration is a day or less. Frue Scirocco events occur infrequently and are normally accompanied by dust and/or "red rain" as wind borne particles from North Africa reach the area.	b. Inbound/outbound vessels should not encounter any significant problems while entering or leaving the Port. Ships inbound to the Italian Navy Base may need extra lines for mooring. Ships intending to anchor outside the outer breakwater should be aware of problems caused by winds from SSE or 5.	b. SE'ly winds at La Spezia are most often caused by low pressure systems located in the Gulf of Benoa, but are infrequently caused by Scirocco events. (1) Benoa lows. The following is an abbreviated listing of various aspects of Genoa low behavior. (a) Cyclogenesis 1. A lee trough often is present in the Gulf of Genoa when a cold or occluded front is moving into western France. The trough remains stationary until arrival of the front, at which time significant cyclogenesis occurs.
			2. A good indication of rapid development of a Genoa low is the appearance of cold air from the NE in the Po Valley. 3. Genoa lows occur almost simultaneously with the onset of a Mistral in the Bulf of Lion, and invariably form when conditions are right for Mistral to occur. (b) Associated weather: Convective activity associated with a Geno low has a periodicity of about 18 hr, starting with the initial cold frontal passage, and is most pronounced with a stationary low. The most intense convective activity occurs at 36 hr intervals. (c) Miscellaneous 1. A residual low pressure trough generally remains over the Bulf of Genoa even after the primary low has moved well out of the region. The trough can remain several days. 2. Centers of Genoa con be poorly organized: Strong pressure gradients frequently are found far from the low's geographic center.
Strongest in Winter	c. SW'ly winds/waves - Any relatively strong	c. Inbound/outbound yessels should experience no significant problems entering	(2) Stirocco events (a) Clouds forming on the SE slope of Mt. Parodi (NW of La Spezia) may indicate the beginning of a Scirocco. (b) Scirocco minds usually will increase slowly, so ample advance warning is provided by the time taken to develop a full Scirocco situation.
Occurs in Spring Scamer and Autumn	c. SM'ly winds/waves - Any relatively strong wind with a direction from SM through MNW has the potential to generate sea/swell that would adversely impact the outer anchorages. Any wind with a strong M component would reach the inner harbor as SM after passing through valleys in peninsula M of the Port. Normally limited to force 5 (17-21 kt) in harbor area. May be caused by low pressure N and/or NE of La Spezia or as post frontal winds following passage of a cold/occluded front.	or leaving Port when in the lee of Palmaria Island and the peninsula which forms the western side of Rada De La Spezia. Once the lee is lost however, vessels are exposed to open-sea conditions. The outer anchorages are exposed, and are not viable alternatives during heavy weather.	c. Cyclogenesis or low pressure center enving over N Italy or Bulf of Venice may result in SM wind at La Spezia. SW to M winds may follow the passage of a cold or occluded front. If high pressure builds in behind a SE'ly moving low pressure center, the winds will likely continue to veer, becoming N as the low moves.

Table 3-2. (Continued)

VESSEL LOCATION/SITUATION	POTENTIAL HAZARD	EFFECT - PRECAUTIONARY/EVASIVE ACTIONS	ADVANCE INDICATORS AND OTHER INFORMATION ABOUT POTENTIAL HAZARD
5. <u>Small boats.</u> Strongest in Winter Occurs in Spring and Autumn Weakest in Summer	a. N'ly winds/waves - Locally called Iramontana. Caused by building high pressure over northern Italy as a low pressure center moves SE from the Bulf of Benga. Maximum winds normally force 6 to 7 (22-33 kt). Raises a N'ly sea to 3 ft in the outer harbor which reflects off the N side of the outer breakwater and creates a chop in the nearby waters.	a. Winds/waves pose no problem in the inner harbor. Since wave heights are limited to about 3 ft in a strong situation, operations in the outer harbor may still be conducted. Caution is necessary in the chop raised by the reflected wind haves near the outer breakwater. BOATING HAS NEVER BEEN CANCELLED INSIDE IHE BREAKWATER DUE TO BAD MEATHER. Boating to/from the breakwater and the outer anchorages may be curtailed in a strong event. Be aware of wind chill factor.	a. N'ly winds are the result of building high pressures over N Italy and a SE'ly moving low pressure center.
Strongest in Winter Occurs in Spring and Autuan Weakest in Summer	b. SE'ly winds - Locally called Scirocco. Caused by low pressure systems developing in or moving into the Bulf of Genoa. Velocities to force 7 to 8 (28-40 kt) can be expected 3-4 times per year. Stronger events may last up to 3 days, but the usual duration is a day or less. True Scirocco events occur infrequently and are normally accompanied by dust and/or 'red rain' as wind borne particles from North Africa reach the area.	b. Winds/waves pose no problem in the harbor due to lack of significant fetch length. BDATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER DUE TO BAD MEATHER. Boating to/from the breakwater and the outer anchorages may be curtailed in a strong event or if the wind is from SSC or S. Local water taxis will not go outside the breakwater in winds of force 6 (22-27 kt) or greater.	b. SE'ly winds at La Spezia are most often caused by low pressure systems located in the Gulf of Benoa, but are infrequently caused by Scirocco events. (1) Genoa lows. The following is an abbreviated listing of various aspects of Benoa low behavior. (a) Cyclogenesis 1. A lee trough often is present in the Bulf of Benoa when a cold or occluded front is moving into western France. The trough remains stationary until arrival of the front, at which time significant cyclogenesis occurs. 2. A good indication of rapid development of a Senoa low is the appearance of cold air from the NC in the Po Valley. 3. Benoa lows occur almost simultaneously with the onset of a Mistral in the Gulf of Lion, and invariably form when conditions are right for a Mistral to occur. (b) Associated weather: Convective activity associated with a Benoa low has a periodicity of about 18 hr, starting with the initial cold frontal passage, and is most promounced with a stationary low. The most intense convective activity occurs at 35 hr intervals. (c) Miscellaneous 1. A residual low pressure trough generally remains over the Bulf of Genoa even after the primary low has moved well out of the region. The trough can remain several days. 2. Centers of Genoa lows can be poorly organized: Strong pressure gradents frequently are found far from the low's geographic center. (2) Scirocco events (3) Clouds forming on the SE slope of Mt. Parodi (NM of La Spezia) may indicate the beginning of a Scirocco. (b) Scirocco events (c) Historian of Benoa lows can be poorly organized: Strong pressure gradents frequently are found far from the low's geographic center. (3) Clouds forming on the SE slope of Mt. Parodi (NM of La Spezia) may indicate the beginning of a Scirocco.
	c. SM'ly winds/waves - Any relatively strong wind with a direction from SM through WNM has the potential to generate sea/swell that would adversely affect boat runs to the outer anchorages. Any wind with a strong W component would reach the inner harbor as SM after passing through valleys in peninsula W of the Port. Normally limited to force 5 (17-21 kt) in harbor area. May be caused by low pressure N and/or NE of La Spezia or as post frontal winds following passage of a cold/occluded front.	c. Doerations inside the outer breakwater are not significantly affected. BOATING HAS NEVER BEEN CANCELLED INSIDE THE BREAKWATER DUE TO BAD WEATHER. Boating to/from the outer anchorages may not be feasible once the lee of Palmaria Island and the peninsula on the W side of Rada de La Spezia is lost.	warning is provided by the time taken to develop a full Scirocco situation. c. Cyclogenesis or low pressure center moving over N Italy or Gulf of Venice may result in SM wind at its Spezia. SM to N winds may follow the passage of cold or occluded front. If high pressure builds in behind a SE'ly moving low pressure center, the winds will likely continue to veer, becoming N as the low moves.

REFERENCES

Brody, L. R. and M. J. R. Nestor, 1980: Regional Forecasting Aids for the Mediterranean Basin, NAVENVPREDRSCHFAC Technical Report TR 80-10. Naval Environmental Prediction Research Facility, Monterey, CA 93941.

Hydrographer of the Navy, 1965: <u>Mediterranean Pilot</u>, Volume II. Hydrographer of the Navy, London, England.

Kotsch, W. J., 1983: <u>Weather for the Mariner</u>, Third Edition. Naval Institute Press, Annapolis, MD.

FICEURLANT, 1987: <u>Port Directory for La Spezia (1984)</u>, <u>Italy</u>. Fleet Intelligence Center Europe and Atlantic, Norfolk, VA.

PORT VISIT INFORMATION

AUGUST 1987. NEPRF meteorologists R. Fett and D. Perryman met with the Port Captain Como Marzaroli and Chief Pilot Mr. Casareto to obtain much of the information included in this port evaluation.

APPENDIX A

General Purpose Oceanographic Information

This section provides general information on wave and wave climatology as used in this study. The forecasting material is not harbor specific. material in paragraphs A.1 and A.2 was extracted from No. 603, Practical Methods for Observing and Pub. Forecasting Ocean Waves (Pierson, Neumann, and James. 1955). The information on fully arisen wave conditions and wave conditions within the fetch region (A.4) (A.3)is based on the JONSWAP model. This model was developed from measurements of wind wave growth over the North Sea in 1973. The JONSWAP model is considered appropriate for enclosed sea where residual wave an activity is minimal and the onset and end of locally forced wind events occur rapidly (Thornton, 1986). where waves are fetch limited and growing (Hasselmann, et al., 1976). Enclosed sea, rapid onset/subsiding local winds, and fetch limited waves are more representative of the Mediterranean waves and winds than the conditions of the North Atlantic from which data was used Pierson and Moskowitz (P-M) Spectra (Neumann and Pierson 1966). The P-M model refined the original spectra of H.O. 603, which over developed wave heights.

The primary difference in the results of the JONSWAP and P-M models is that it takes the JONSWAP model longer to reach a given height or fully developed seas. In part this reflects the different starting wave conditions. Because the propagation of waves from surrounding areas into semi-enclosed seas, bays, harbors, etc. is limited, there is little residual wave action following periods of locally light/calm winds and the sea surface is nearly flat. A local wind developed wave growth is therefore slower than wave growth in the open ocean where some residual wave action is generally always

present. This slower wave development is a built in bias in the formulation of the JONSWAP model which is based on data collected in an enclosed sea.

A.1 Definitions

Waves that are being generated by local winds are called "SEA". Waves that have traveled out of the generating area are known as "SWELL". Seas are chaotic in period, height and direction while swell approaches a simple sine wave pattern as its distance from the generating area increases. An in-between state exists for a few hundred miles outside the generating area and is a condition that reflects parts of both of the above definitions. In the Mediterranean area, because its fetches and open sea expanses are limited, SEA or IN- BETWEEN conditions will prevail. The "SIGNIFICANT WAVE HEIGHT" is defined as the average value of the heights of the one-third highest waves. PERIOD and WAVE LENGTH refer to the time between passage of, and distances between, two successive crests on the sea surface. The FREQUENCY is the reciprocal of the period (f = 1/T) therefore as the period increases the frequency decreases. Waves result from the transfer of energy from the wind to the sea surface. The area over which the wind blows is known as the FETCH, and the length of time that the wind has blown is the DURATION. The characteristics of waves (height, length, and period) depend on the duration, fetch, velocity of the wind. There is a continuous generation of small short waves from the time the wind starts until it stops. With continual transfer of energy from the wind to the sea surface the waves grow with the older waves leading the growth and spreading the energy over a greater range of frequencies. Throughout the growth cycle a SPECTRUM of ocean waves is being developed.

A.2 Wave Spectrum

Wave characteristics are best described by means of their range of frequencies and directions or their spectrum and the shape of the spectrum. If the spectrum of the waves covers a wide range of frequencies directions (known as short-crested conditions), SEA conditions prevail. If the spectrum covers a narrow range frequencies and directions (long crested conditions), SWELL conditions prevail. The wave spectrum depends on the duration of the wind, length of the fetch, and on the wind velocity. At a given wind speed given state of wave development, each spectrum has a band frequencies where most of the total energy is concentrated. As the wind speed increases the range of significant frequencies extends more and more toward lower frequencies (longer periods). The frequency of maximum energy is given in equation 1.1 where v is the wind speed in knots.

$$f_{max} = \frac{2.476}{V}$$
 (1.1)

The wave energy, being a function of height squared, increases rapidly as the wind speed increases and the maximum energy band shifts to lower frequencies. This results in the new developing smaller waves (higher frequencies) becoming less significant in the energy spectrum as well as to the observer. As larger waves develop an observer will pay less and less attention to the small waves. At the low frequency (high period) end the energy drops off rapidly, the longest waves are relatively low and extremely flat, and therefore also masked by the high energy frequencies. The result is that 5% of the upper frequencies and 3% of the lower frequencies can be cut-off and only the remaining

frequencies are considered as the "significant part of the wave spectrum". The resulting range of significant frequencies or periods are used in defining a fully arisen sea. For a fully arisen sea the approximate average period for a given wind speed can be determined from equation (1.2).

$$T = 0.285v$$
 (1.2)

Where v is wind speed in knots and T is period in seconds. The approximate average wave length in a fully arisen sea is given by equation (1.3).

$$\bar{L} = 3.41 \, \bar{T}^2$$
 (1.3)

Where \overline{L} is average wave length in feet and \overline{T} is average period in seconds.

The approximate average wave length of a fully arisen sea can also be expressed as:

$$L = .67$$
"L" (1.4)

where "L" = $5.12T^2$, the wave length for the classic sine wave.

A.3 Fully Arisen Sea Conditions

For each wind speed there are minimum fetch (n mi) and duration (hr) values required for a fully arisen sea to exist. Table A-1 lists minimum fetch and duration values for selected wind speeds, values of significant wave (average of the highest 1/3 waves) period and height, and wave length of the average wave during developing and fully arisen seas. The minimum duration time assumes a start from a flat sea. When pre-existing

lower waves exist the time to fetch limited height will be shorter. Therefore the table duration time represents the maximum duration required.

Table A-1. Fully Arisen Deep Water Sea Conditions Based on the JONSWAP Model.

;	Wind Speed	;	Minimu Fetch/		ration	;			(H1/3) Height	1	Wave Len Developi		
;	(kt)	,	(n mi)		(hrs)	ļ	(sec)	(ft)	1		_	Arisen !
1		1				. !				1	L X (.5)		X (.67):
;	10	1	28 /	,	4	1	4	1	2	;	41	/	55 ;
;	15	1	55 /	,	6	ł	6	1	4	1	92	1	123
;	20	;	110 /	′	8	1	8	1	8	;	164	/	220
1	25	- }	160 /	,	11	ļ	9	1	12	1	208	/	27B ·
1	30	- [210 /	,	13	į	11	1	16		310	/	415
1	35	1	310 /	,	15	1	13	1	22	i	433	,	580
1	40	-	410 /	′	17	1	15	1	30	i	576	1	772

NOTES:

- Depths throughout fetch and travel zone must be greater than 1/2 the wave length, otherwise shoaling and refraction take place and the deep water characteristics of waves are modified.
- For the classic sine wave the wave length (L) equals 5.12 times the period (T) squared (L = 5.12T²). As waves develop and mature to fully developed waves and then propagate out of the fetch area as swell their wave lengths approach the classic sine wave length. Therefore the wave lengths of developing waves are less than those of fully developed waves which in turn are less than the length of the resulting swell. The factor of .5 (developing) and .67 (fully developed) reflect this relationship.

A.4 Wave Conditions Within The Fetch Region

Waves produced by local winds are referred to as <u>SEA</u>. In harbors the local sea or wind waves may create hazardous conditions for certain operations. Generally within harbors the fetch lengths will be short and therefore the growth of local wind waves will be fetch limited. This implies that there are locally determined upper limits of wave height and period for each wind velocity. Significant changes in speed or direction will result in generation of a new wave group with a new set of height and period limits. Once a fetch limited sea reaches its upper limits no further growth will occur unless the wind speed increases.

Table A-2 provides upper limits of period and height for given wind speeds over some selected fetch lengths. The duration in hours required to reach these upper limits (assuming a start from calm and flat sea conditions) is also provided for each combination of fetch length and wind speed. Some possible uses of Table A-2 information are:

- If the only waves in the area are locally generated wind waves, the Table can be used to forecast the upper limit of sea conditions for combinations of given wind speeds and fetch length.
- 2) If deep water swell is influencing the local area in addition to locally generated wind waves, then the Table can be used to determine the wind waves that will combine with the swell. Shallow water swell conditions are influenced by local bathymetry (refraction and shoaling) and will be addressed in each specific harbor study.
- 3) Given a wind speed over a known fetch length the maximum significant wave conditions and time needed to reach this condition can be determined.

Table A-2. Fetch Limited Wind Wave Conditions and Time Required to Reach These Limits (Based on JONSWAP Model). Enter the table with wind speed and fetch length to determine the significant wave height and period, and time duration needed for wind waves to reach these limiting factors. All of the fetch/speed combinations are fetch limited except the 100 n mi fetch and 18 kt speed.

Format: height (feet)/period (seconds)
duration required (hours)

Fetch \	Wind Speed				:
! Length	\ 18	24	30 1	36	42
(n mi)	;		<u> </u>		1
10	 2/3-4 1-2	 3/3-4 2	3-4/4	4/4-5 1-2	5/5 1-2
20	 3/4-5 2-3	 4/4-5 3		6/5-6 3-4	7/5-6 3
30	 3-4/5 3	: : 5/5-6 : 4	6/6 6/6 3-4	7/6 3-4	B/6-7 3
1 40	 4-5/5-6 4-5	l l 5/6 l 4		8/7 4	9-10/7-8 3-4
100	 5/6-7 ¹ 5-6	! ! 9/8 ! 8		13/9 7	 15-16/9-10 7

¹⁸ kt winds are not fetch limited over a 100 n mi fetch.

An example of expected wave conditions based on Table A-2 follows:

WIND FORECAST OR CONDITION

An offshore wind of about 24 kt with a fetch limit of 20 n mi (ship is 20 n mi from the coast) is forecast or has been occurring.

SEA FORECAST OR CONDITION

From Table A-2: If the wind condition is forecast to last, or has been occurring, for at least 3 hours:

Expect sea conditions of 4 feet at 4-5 second period to develop or exist. If the condition lasts less than 3 hours the seas will be lower. If the condition lasts beyond 3 hours the sea will not grow beyond that developed at the end of about 3 hours unless there is an increase in wind speed or a change in the direction that results in a longer fetch.

A.5 Wave Climatology

The wave climatology used in these harbor studies is based on 11 years of Mediterranean SOWM output. MED-SOWM is discussed in Volume II of the U.S. Oceanography Command Numerical Environmental Products Manual (1986). A deep water MED-SOWM grid point selected as representative of the deep water wave conditions outside each harbor. The deep water waves were then propagated into the shallow water areas. linear wave theory and wave refraction computations the shallow water climatology was derived from the modified deep water wave conditions. This climatology does not include the local wind generated seas. This omission, by design, is accounted for by removing all wave data for periods less than 6 seconds in the climatology. These shorter period waves are typically dominated by locally generated wind waves.

A.6 Propagation of Deep Water Swell Into Shallow Water Areas

When deep water swell moves into shallow water the wave patterns are modified, i.e., the wave heights and directions typically change, but the wave period remains constant. Several changes may take place including shoaling as the wave feels the ocean bottom, refraction as the wave crest adjusts to the bathymetry pattern, changing so that the crest becomes more parallel to the bathymetry contours, friction with the bottom sediments, interaction with currents, and adjustments caused by water temperature gradients. In this work. only shoaling and refraction effects are considered. Consideration of the other factors are beyond resources available for this study and, furthermore, they are considered less significant in the harbors of this study than the refraction and shoaling factors.

To determine the conditions of the deep water waves in the shallow water areas the deep water

conditions were first obtained from the Navy's operational MED-SOWM wave model. The bathymetry for the harbor/area of interest was extracted from available charts and digitized for computer use. Figure A-1 is a sample plot of bathymetry as used in this project. path refraction/shoaling program was run for selected combinations of deep water wave direction and period. The selection was based on the near deep water climatology and harbor exposure. Each study requires a number of ray path computations. Typically there are 3 or 4 directions (at 30° increments) and 5 or 6 periods (at 2 second intervals) of concern for each study. This results in 15 to 24 plots per area/harbor. To reduce this to a manageable format for quick reference, specific locations within each study area were selected and the information was summarized and is presented in the specific harbor studies in tabular form.




Figure A-1. Example plot of bathymetry (Naples harbor) as used in this project. For plotting purposes only, contours are at 50 fathom intervals from an initial 10 fathoms to 110 fathoms, and at 100 fathom intervals thereafter. The larger size numbers identify specific anchorage areas addressed in the harbor study.

REFERENCES

Hasselmann, K. D., D. B. Ross, P. Muller, and W. Sell, 1976: A parametric wave prediction model. <u>J. Physical Oceanography</u>, Vol. 6, pp. 208-228.

Neumann, G., and W. J. Pierson Jr., 1966: <u>Principles of</u>
<u>Physical Oceanography</u>. Prentice-Hall, Englewood Cliffs.

Pierson, W. J. Jr., G. Neumann, and R. W. James, 1955: Practical Methods for Observing and Forecasting Ocean Waves, H. D. Pub. No. 603.

Thornton, E. B., 1986: <u>Unpublished lecture notes for DC 3610</u>, <u>Waves and Surf Forecasting</u>. Naval Postgraduate School, Monterey, CA.

U. S. Naval Oceanography Command, 1986: Vol. II of the
U. S. Naval Oceanography Command Numerical Environmental
Products Manual.

DISTRIBUTION LIST

SNDL		
21A1 21A3 22A1 22A3	CINCLANTFLT CINCUSNAVEUR COMSECONDFLT COMSIXTHFLT	
23B3 24A1 24D1 24E 24G1	Special Force Commander EUR Naval Air Force Commander LANT Surface Force Commander LANT Mine Warfare Command Submarine Force Commander LANT	
26QQ1 28A1 28B1 28D1	Special Warfare Group LANT Carrier Group LANT (2) Cruiser-Destroyer Group LANT (2) Destroyer Squadron LANT (2)	
28J1 28K1 28L1 29A1 29B1	Service Group and Squadron LANT (2) Submarine Group and Squadron LANT Amphibious Squadron LANT (2) Guided Missile Cruiser LANT Aircraft Carrier LANT	
29D1 29E1 29F1 29G1	Destroyer LANT (DD 931/945 Class) Destroyer LANT (DD 963 Class) Guided Missile Destroyer LANT Guided Missile Frigate (LANT)	
2911 29J1 29K1 29L1 29N1	Frigate LANT (FF 1098) Frigate LANT (FF 1040/1051 Class) Frigate LANT (FF 1052/1077 Class) Frigate LANT (FF 1078/1097 Class) Submarine LANT (SSN)	
29Q 29R1 29AA1 29BB1	Submarine LANT SSBN Battleship Lant (2) Guided Missile Frigate LANT (FFG 7) Guided Missile Destroyer (DDG 993)	
31A1 31B1 31G1 31H1 31I1	Amphibious Command Ship LANT (2) Amphibious Cargo Ship LANT Amphibious Transport Ship LANT Amphibious Assault Ship LANT (2) Dock Landing Ship LANT	
31J1 31M1 32A1 32C1	Dock Landing Ship LANT Tank Landing Ship LANT Destroyer Tender LANT Ammunition Ship LANT	
32G1 32H1 32N1 32Q1 32S1	Combat Store Ship LANT Fast Combat Support Ship LANT Oiler LANT Replenishment Oiler LANT Repair Ship LANT	
32X1 32DD1 32EE1 32KK	Salvage Ship LANT Submarine Tender LANT Submarine Rescue Ship LANT Miscellaneous Command Ship	
32QQ1 32TT	Salvage and Rescue Ship LANT Auxiliary Aircraft Landing Training	Ship

42N1 42P1	Air Anti-Submarine Squadron VS LANT Patrol Wing and Squadron LANT
42BB1	Helicopter Anti-Submarine Squadron HS LANT
	mericopter Anti-Submarine Squadron HS LANT
42CC1	Helicopter Anti-Submarine Squadron Light HSL LANT
C40	Monterey, Naples, Sigonella and Souda Bay only
FD2	Oceanographic Office - COMNAVOCEANCOM
FD3	Fleet Numerical Oceanography Center - FNOC
FD4	Oceanography Center - NAVEASTOCEANCEN
FD5	Oceanography Command Center - COMNAVOCEANCOM

copy to:

21A2	CINCPACFLT
22A2	Fleet Commander PAC
24F	Logistics Command
24H1	Fleet Training Command LANT
28A2	Carrier Group PAC (2)
29B2	Aircraft Carrier PAC (2)
29R2	Battleships PAC (2)
31A2	Amphibious Command Ship PAC (2)
31H2	Amphibious Assault Ship PAC (2)
FA2	Fleet Intelligence Center
FC14	Air Station NAVEUR
FDl	Oceanography Command
USDAO	France, Israel, Italy and Spain

Stocked:

NAVPUBFORMCEN (50 copies)

NAVENVPREDRSCHFAC SUPPLEMENTARY DISTRIBUTION

COMMANDER IN CHIEF USCINCLANT COMMANDING GENERAL (G4) U.S. CENTRAL COMMAND NAVAL BASE FLEET MARINE FORCE, ATLANTIC NORFOLK, VA 23511 MACDILL AFB, FL 33608 ATTN: NSAP SCIENCE ADVISOR NORFOLK, VA 23511 ASST. FOR ENV. SCIENCES CHIEF OF NAVAL RESEARCH (2) USCINCENT LIBRARY SERVICES, CODE 784 ASST. SEC. OF THE NAVY (R&D) ATTN: WEATHER DIV. (CCJ3-W) ROOM 5E731, THE PENTAGON WASHINGTON, DC 20350 BALLSTON TOWER #1 MACDILL AFB, FL 33608-7001 800 QUINCY ST. ARLINGTON, VA 22217-5000 OFFICE OF NAVAL RESEARCH OFFICE OF NAVAL RESEARCH OFFICE OF NAVAL RESEARCH ATTN: PROGRAM MANAGER, 1122CS ENV. SCI. PROGRAM, CODE 112 CODE 1122AT, ATMOS. SCIENCES ARLINGTON, VA 22217-5000 ARLINGTON, VA 22217-5000 ARLINGTON, VA 22217-5000 OFFICE OF NAVAL RESEARCH OFFICE OF NAVAL RESEARCH OFFICE OF NAVAL RESEARCH ATTN: HEAD, OCEAN SCIENCES DIV CODE 1122 PO, PHYSICAL OCEANO. CODE 1122 MM, MARINE METEO. ARLINGTON, VA 22217-5000 ARLINGTON, VA 22217-5000 CODE 1122 ARLINGTON, VA 22217-5000 CHIEF OF NAVAL OPERATIONS CHIEF OF NAVAL OPERATIONS OFFICE OF NAVAL TECHNOLOGY NAVY DEPT., OP-622C (0P-006)ONR (CODE 22) U.S. NAVAL OBSERVATORY WASHINGTON, DC 20350 800 N. QUINCY ST. ARLINGTON, VA 22217-5000 WASHINGTON, DC 20390 CHIEF OF NAVAL OPERATIONS CHIEF OF NAVAL OPERATIONS CHIEF OF NAVAL OPERATIONS U.S. NAVAL OBSERVATORY OP-952D NAVY DEPT. OP-986G U.S. NAVAL OBSERVATORY DR. RECHNITZER, OP-952F WASHINGTON, DC 20350 WASHINGTON, DC 20390 34TH & MASS AVE. WASHINGTON, DC 20390 COMMANDANT OF THE MARINE CORPS DIRECTOR CHIEF OF NAVAL OPERATIONS NATIONAL SECURITY AGENCY HDQ, U.S. MARINE CORPS OP-953 ATTN: LIBRARY (2CO29) WASHINGTON, DC 20380 NAVY DEPARTMENT FT. MEADE, MD 20755 WASHINGTON, DC 20350

CHARLESTON, SC 29408-6475

OFFICER IN CHARGE

NAVOCEANCOMDET

NAVAL STATION

OJCS/J3/ESD

THE PENTAGON, ROOM 2B887

WASHINGTON, DC 20301-5000

OFFICER IN CHARGE

BOX 16

U.S. NAVOCEANCOMDET

FPO NEW YORK 09593-5000

OFFICER IN CHARGE NAVOCEANCOMDET NAVAL EDUCATION & TRNG CENTER NEWPORT, RI 02841-5000

OFFICER IN CHARGE U.S. NAVOCEANCOMDET APO NEW YORK 09406-5000

COMMANDING OFFICER NAVAL RESEARCH LAB ATTN: LIBRARY, CODE 2620 WASHINGTON, DC 20390

OFFICE OF NAVAL RESEARCH SCRIPPS INSTITUTION OF **OCEANOGRAPHY** LA JOLLA, CA 92037

COMMANDING OFFICER NAVAL OCEAN RSCH & DEV ACT NSTL, MS 39529-5004

COMMANDING OFFICER FLEET INTELLIGENCE CENTER (EUROPE & ATLANTIC) NORFOLK, VA 23511

COMMANDER NAVAL OCEANOGRAPHY COMMAND NSTL, MS 39529-5000

COMNAVOCEANCOM ATTN: CODE N5 NSTL, MS 39529-5000 SUPERINTENDENT LIBRARY REPORTS U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402

CHAIRMAN OCEANOGRAPHY DEPT. U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402 DIRECTOR OF RESEARCH U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402

NAVAL POSTGRADUATE SCHOOL OCEANOGRAPHY DEPT. MONTEREY, CA 93943-5000

LIBRARY NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943-5002

PRESIDENT NAVAL WAR COLLEGE GEOPHYS. OFFICER, NAVOPS DEPT. NAVAL AIR STATION NEWPORT, RI 02841

COMMANDER NAVAL SAFETY CENTER NORFOLK, VA 23511

COMSPAWARSYSCOM ATTN: CAPT. R. PLANTE CODE 3213, NAVY DEPT. WASHINGTON, DC 20363-5100

COMMANDER, D.W. TAYLOR NAVAL SHIP RSCH. & DEV. CENTER SURFACE SHIP DYNAMICS BRANCH ATTN: S. BALES BETHESDA, MD 20084-5000

COMMANDER NAVSURFWEACEN, CODE R42 DR. B. KATZ, WHITE OAKS LAB SILVER SPRING, MD 20903-5000

DIRECTOR NAVSURFWEACEN, WHITE OAKS NAVY SCIENCE ASSIST. PROGRAM SILVER SPRING, MD 20903-5000 COMMANDING GENERAL FLEET MARINE FORCE, LANT (G4) ATTN: NSAP SCIENCE ADVISOR NORFOLK, VA 23511

USAFETAC/TS SCOTT AFB, IL 62225

3350TH TECH. TRNG GROUP TTGU/2/STOP 623 CHANUTE AFB, IL 61868

OFFICER IN CHARGE SERVICE SCHOOL COMMAND DET. CHANUTE/STOP 62 CHANUTE AFB, IL 61868

COMMANDING OFFICER U.S. ARMY RESEARCH OFFICE ATTN: GEOPHYSICS DIV. P.O. BOX 12211 RESEARCH TRIANGLE PARK, NC 27709

COMMANDER
COASTAL ENGINEERING RSCH CEN
KINGMAN BLDG.
FT. BELVOIR, VA 22060

DIRECTOR
LIBRARY, TECH. INFO. CEN.
ARMY ENG. WATERWAYS STN.
VICKSBURG, MS 39180

DIRECTOR (12)
DEFENSE TECH. INFORMATION
CENTER, CAMERON STATION
ALEXANDRIA, VA 22314

DIRECTOR, ENV. & LIFE SCI.
OFFICE OF UNDERSECRETARY OF
DEFENSE FOR RSCH & ENG E&LS
RM. 3D129, THE PENTAGON
WASHINGTON, DC 20505

CENTRAL INTELLIGENCE AGENCY ATTN: OCR STANDARD DIST. WASHINGTON, DC 20505 DIRECTOR, TECH. INFORMATION DEFENSE ADV. RSCH PROJECTS 1400 WILSON BLVD. ARLINGTON, VA 22209

COMMANDANT
DEFENSE LOGISTICS STUDIES
INFORMATION EXCHANGE
ARMY LOGISTICS MANAGEMENT
CENTER
FORT LEE, VA 23801

COMMANDANT U.S. COAST GUARD WASHINGTON, DC 20226 CHIEF, MARINE SCI. SECTION U.S. COAST GUARD ACADEMY NEW LONDON, CT 06320

COMMANDING OFFICER USCG RESTRACEN YORKTOWN, VA 23690

COMMANDING OFFICER
USCG RSCH & DEV. CENTER
GROTON, CT 06340

OCEANOGRAPHIC SERVICES DIV. NOAA 6010 EXECUTIVE BLVD. ROCKVILLE, MD 20852

FEDERAL COORD. FOR METEORO. SERVS. & SUP. RSCH. (OFCM) 11426 ROCKVILLE PIKE SUITE 300 ROCKVILLE, MD 20852 NATIONAL CLIMATIC CENTER ATTN: L. PRESTON D542X2 FEDERAL BLDG. - LIBRARY ASHEVILLE. NC 28801 DIRECTOR NATIONAL OCEANO. DATA CENTER E/OC23, NOAA WASHINGTON, DC 20235

NOAA RSCH FACILITIES CENTER P.O. BOX 520197 MIAMI, FL 33152 DIRECTOR
ATLANTIC MARINE CENTER
COAST & GEODETIC SURVEY, NOAA
439 W. YORK ST.
NORFOLK, VA 23510

CHIEF, INTERNATIONAL AFFAIRS NATIONAL WEATHER SERVICE 8060 13TH STREET SILVER SPRING, MD 20910

HEAD
OFFICE OF OCEANO. & LIMNOLOGY
SMITHSONIAN INSTITUTION
WASHINGTON, DC 20560

SCRIPPS INSTITUTION OF OCEANOGRAPHY, LIBRARY DOCUMENTS/REPORTS SECTION LA JOLLA, CA 92037

WOODS HOLE OCEANO. INST. DOCUMENT LIBRARY LO-206 WOODS HOLE, MA 02543

SCIENCE APPLICATIONS
INTERNATIONAL CORP. (SAIC)
205 MONTECITO AVE.
MONTEREY, CA 93940

OCEANROUTES, INC. 680 W. MAUDE AVE. SUNNYVALE, CA 94086-3518 MR. W. G. SCHRAMM/WWW
WORLD METEOROLOGICAL
ORGANIZATION
CASE POSTALE #5, CH-1211
GENEVA, SWITZERLAND

DIRECTOR, INSTITUTE OF PHYSICAL OCEANOGRAPHY HARALDSGADE 6 2200 COPENHAGEN N. DENMARK

MINISTRY OF DEFENCE NAVY DEPARTMENT ADMIRALTY RESEARCH LAB TEDDINGTON, MIDDX ENGLAND

METEOROLOGIE NATIONALE SMM/DOCUMENTATION 2, AVENUE RAPP 75340 PARIS CEDEX 07 FRANCE

DIRECTION DE LA METEOROLOGIE ATTN: J. DETTWILLER, MN/RE 77 RUE DE SEVRES 92106 BOULOGNE-BILLANCOURT CEDEX, FRANCE

INSTITUT FUR MEERESKUNDE DER UNIVERSITAT HAMBURG HEIMHUDERSTRASSE 71 2000 HAMBURG 13 FEDERAL REPUBLIC OF GERMANY

CONSIGLIO NAZIONALE DELLE RICERCHE ISTITUTO TALASSOGRAFICO DI TRIESTE, VIALE R. GESSI 2 34123 TRIESTE, ITALY DIRECTOR OF NAVAL
OCEANO. & METEOROLOGY
MINISTRY OF DEFENCE
OLD WAR OFFICE BLDG.
LONDON, S.W.1. ENGLAND

COMMANDER IN CHIEF FLEET ATTN: STAFF METEOROLOGIST & OCEANOGRAPHY OFFICER NORTHWOOD, MIDDLESEX HA6 3HP ENGLAND

SERVICE HYDROGRAPHIQUE ET OCEANOGRAPHIQUE DE LA MARINE ESTABLISSEMENT PRINCIPAL RUE DU CHATELLIER, B.P. 426 29275 - BREST CEDEX, FRANCE

OZEANOGRAPHISCHE FORSCHUNGSANTALT BUNDESWEHR LORNSENSTRASSE 7, KIEL FEDERAL REPUBLIC OF GERMANY

DIRECTOR, DEUTSCHES
HYDROGRAPHISCHES INSTITUT
TAUSCHSTELLE, POSTFACH 220
D2000 HAMBURG 4
FEDERAL REPUBLIC OF GERMANY

DIRECTOR, SACLANT ASW RESEARCH CENTRE VIALE SAN BARTOLOMEO, 400 I-19026 LA SPEZIA, ITALY THE BRITISH LIBRARY
SCIENCE REFERENCE LIBRARY (A)
25 SOUTHAMPTON BLDGS.
CHANCERY LANE
LONDON WC2A 1AW

LIBRARY, INSTITUTE OF OCEANOGRAPHIC SCIENCES ATTN: DIRECTOR WORMLEY, GODALMING SURRY GUB 5UB, ENGLAND

METEOROLOGIE NATIONALE 1 QUAI BRANLY 75, PARIS (7) FRANCE

INSTITUT FUR MEERESKUNDE AN DER UNIVERSITAT KIEL DUSTERNBROOKER WEG 20 23 KIEL FEDERAL REPUBLIC OF GERMANY

ISTITUTO UNIVERSITARIO NAVALE FACILTA DI SCIENZE NAUTICHE ISTITUTO DI METEOROLOGIA E OCEANOGRAFIA, 80133 NAPOLI -VIA AMM, ACTON, 38 ITALY DUDLEY KNOX LIBRARY - RESEARCH REPORTS

U236856